

**Antriscus cerefolium (L.) Hoffm.
National protocol: TP/KRV/2**

Examination office :	Naktuinbouw	
Reference of the protocol :	TP/KRV/2	
Date of preparation of the protocol :	15-12-2020	
Date of entry into force of the protocol :	15/12/2020	
Botanical taxon:	Antriscus cerefolium (L.) Hoffm.	
Common Name (when known):	Chevril/Kervel	
Way of propagation of the plants to be examined :	Self or cross pollinated seed propagated <input type="checkbox"/> Vegetatively propagated <input type="checkbox"/>	
Number of growing cycles: 1. Vegetatively propagated 1 (1 year) 2 Seed propagated 2 (2 years)	1 <input type="checkbox"/> 2 <input type="checkbox"/> Other <input type="checkbox"/> specify	
List of grouping characteristics :	Yes <input type="checkbox"/> if yes put as annex I No <input checked="" type="checkbox"/>	
Minimum number of plants in trial :	Vegetative:20	Seed:100
Minimum number of plants observed by measuring or counting:	Vegetative:20	Seed: 25
Give description of when observations should take place :	Observation on the leaves should be made at fully developed plants. Observation on bolting tendency should be made 4 weeks after the plant is fully developed in the summer. (Antriscus is a long-day crop)	
Uniformity: <ul style="list-style-type: none"> - For the assessment of uniformity of vegetatively propagated, self-pollinated seed propagated varieties or F1-hybrids, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 100 plants, 4 off-types are allowed. - For the assessment of uniformity for cross-pollinated varieties, the recommendations for cross-pollinated varieties in the General introduction of UPOV should be applied. The variability within the variety should not exceed the variability of comparable varieties already known. 		
Table of characteristics :	Present <input checked="" type="checkbox"/> Not available <input type="checkbox"/>	

Literature :
(when present, please annex to this document)

Present

Absent

Annex I: Table of characteristics

1 Table of Characteristics

	English	Example Varieties	Note
1.	Plant: height		
	short	Fijne Krul	1
	medium		3
	tall	Donkergroene Laatdoorschietende	5
2.	Plant: width		
	narrow	Fijne Krul	1
	medium	Donkergroene Laatdoorschietende	3
	broad		5
3.	Plant: density of foliage		
	thin		1
	medium	Donkergroene Laatdoorschietende	3
	thick		5
4.	Leaf: attitude		
	erect		1
	semi-erect	Donkergroene Laatdoorschietende	3
	prostate		5
5.	Leaf blade: profile		
	concave	Fijne Krul	1
	straight	Donkergroene Laatdoorschietende	2
	convex	Vertissimo	3
6.	Leaf blade: curling		
	absent or very weak		1
	medium	Donkergroene Laatdoorschietende	2
	strong	Fijne Krul	3
7.	Leaf blade: length		
	short	Fijne Krul	1
	medium	Vertissimo	3
	long		5
8.	Leaf blade: width		
	narrow	Fijne Krul	1
	medium	Vertissimo	3
	broad		5

9.	Leaf blade: intensity of green colour		
	light	Fijne Krul	1
	medium	Donkergroene Laatdoorschietende	3
	dark	Vertissimo	5
10.	Leaf blade: distance between 1st and 2nd pair of leaflets		
	short	Fijne Krul	1
	medium	Donkergroene Laatdoorschietende	3
	long		5
11.	Leaf blade margin: depth of incisions		
	absent or shallow	Fijne Krul	1
	medium	Donkergroene Laatdoorschietende	3
	deep	Vertissimo	5
12.	Leaf blade: spacing of lobes		
	not touching	Vertissimo	1
	touching	Donkergroene Laatdoorschietende	2
	overlapping	Fijne Krul	3
13.	Petiole: length		
	short	Fijne krul	1
	medium		3
	long	Vertissimo	5
14.	Bolting tendency		
	absent or very weak	Donkergroene Laatdoorschietende	1
	medium		2
	strong		3

Annex II: Literature

[Anthriscus cerefolium \(PROSEA\) - PlantUse English \(plantnet-project.org\)](https://plantnet-project.org/)