

Viburnum Simplified standard protocol: SSP/SBA/4

| Examination office: | Naktuinbouw | |
|---|--|---------|
| Reference of the protocol: | SSP/SBA/4 | |
| Date of preparation of the protocol: | 26/08/2022 | |
| Date of entry into force of the protocol: | 26/08/2022 | |
| Botanical taxon: | Viburnum L. Viburnum cassinoides L. Viburnum odoratissimum Ker Gawl. Viburnum opulus L. Viburnum plicatum Thunb. Viburnum rhytidophyllum Hemsl. Viburnum tinus L. | |
| Common Name (when known): | Snowball Tree | |
| Way of propagation of the plants to be examined: | Self or cross pollinated seed propagated □ Vegetatively propagated ⊠ | |
| Number of growing cycles: | 1 ⊠ 2 □ Other □ specify | |
| List of grouping characteristics: | Yes □ if yes put as annex No ☒ | |
| Minimum number of plants in trial: | Vegetative:8 | Seed: - |
| Minimum number of plants observed by measuring or counting: | Vegetative:1 | Seed: - |
| Give description of when observations should take place: | Observation on the flower should take place: at full flowering Observation on the young leaf should take place: in spring Observation on the leaf should take place: at full flowering | |



| | Observation on the berry should take place: at full maturity | | | |
|---|--|--|--|--|
| | Other observations should take place: at full flowering | | | |
| Uniformity: | | | | |
| varieties or F1-hybrids, a population standard of 1% a | For the assessment of uniformity of vegetatively propagated, self-pollinated seed propagated varieties or F1-hybrids, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 10 plants, 1 off-types are allowed. | | | |
| For the assessment of uniformity for cross-pollinated varieties, the recommendations fo cross-pollinated varieties in the General introduction of UPOV should be applied. The variability within the variety should not exceed the variability of comparable varieties already known. | | | | |
| | Present ⊠ | | | |
| Table of characteristics: | Not available □ | | | |
| Literature: | Present ⊠ | | | |
| (when present, please annex to this document) | Absent □ | | | |



Table of characteristics: Viburnum

| | BL 1 III | |
|-------------------|--|--|
| 1. | Plant: growth habit | |
| | Plant: height | |
| 3. | Plant: color of branches | |
| 4. | Plant: color of bark | 2112.0.1 |
| 5. | Young leaf blade: color of upper side | RHS Colour Chart (indicate reference number) |
| 6. | Young leaf blade: color of lower side | |
| 7. | Petiole: length | |
| 8. | Petiole: color | |
| 9. | Leaf blade: length | |
| | Leaf blade: width | |
| | Leaf blade: shape | |
| | Leaf blade: shape of base | |
| | Leaf blade: shape of apex | |
| | Leaf blade: color of upper side | RHS Colour Chart (indicate reference number) |
| 15. | Leaf blade: intensity of anthocyanin coloration | |
| | of upper side | |
| 16. | Leaf blade: color of main vein | |
| 17. | Leaf blade: color of lower side | |
| 18. | Leaf blade: number of incision of margin | |
| 19. | Leaf blade: undulation of margin | |
| 20. | Leaf blade: shape in cross section | |
| 21. | Leaf blade: curvature of longitudinal axis | |
| 22. | Inflorescence: shape | |
| 23. | Inflorescence: height | |
| 24. | Inflorescence: width | |
| 25. | Inflorescence: conspicuousness of fertile flowers | |
| | Flower bud: color | |
| 27. | Flower bud: intensity of anthocyanin coloration | |
| | Sterile flower: diameter of calyx | |
| 29. | Sterile flower: attitude of sepals | |
| | Sterile flower: shape of apex of sepals | |
| | Sterile flower: shape of sepals in cross section | |
| | Sterile flower: main color of inner side of sepals | RHS Colour Chart (indicate reference number) |
| | Sterile flower: secondary color of inner side of | |
| | sepals | |
| 34. | Sterile flower: distribution of secondary color of | |
| | inner side of sepals | |
| 35. | | |
| | Fertile flower: diameter of corolla | |
| 37. | | |
| | | |
| | | |
| | | RHS Colour Chart (indicate reference number) |
| 38. 39. 40. | | RHS Colour Chart (indicate reference number) |



| 41. | Fertile flower: secondary color of inner side of | |
|-----|--|--|
| | petals | |

- 42. Fertile flower: distribution of secondary color of inner side of petals
- 43. Filament: length
- 44. Filament: color
- 45. Berry: diameter
- 46. Berry: shape
- 47. Berry: color

RHS Colour Chart (indicate reference number)

Literature:

The Cambridge Illustrated Glossary of Botanical Terms: by Michael Hickey and Clive King

Name that flower: by Ian Clarke and Heleen Lee Botanisch woordenboek: by Henk Eggelte