

Olearia <u>Simplified standard protocol: SSP/OLE/1</u>

| Examination office: | Naktuinbouw | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------|---------|
| Reference of the protocol: | SSP/OLE/1 | |
| Date of preparation of the protocol: | 11/08/2022 | |
| Date of entry into force of the protocol: | 01/04/2023 | |
| Botanical taxon: | Olearia phlogopappa (Labill.) DC. | |
| Common Name (when known): | Tasmanian daisybush | |
| Way of propagation of the plants to be examined: | Self or cross pollinated seed propagated □ Vegetatively propagated ⊠ | |
| Number of growing cycles: | 1 ⋈ 2 □ Other □ specify Click or tap here to enter text. | |
| List of grouping characteristics: | Yes □ if yes put as annex No ⊠ | |
| Minimum number of plants in trial: | Vegetative:20 | Seed: - |
| Minimum number of plants observed by measuring or counting: | Vegetative:1 | Seed: - |
| Give description of when observations should take place: | - | |
| Uniformity: For the assessment of uniformity of vegetatively propagated, self-pollinated seed propagated varieties or F1-hybrids, a population standard of 1 % and an acceptance probability of at least 95% should be applied. In the case of a sample size of 20 plants, 1 off-types is allowed. For the assessment of uniformity for cross-pollinated varieties, the recommendations for cross-pollinated varieties in the General introduction of UPOV should be applied. The variability within the variety should not exceed the variability of comparable varieties already known. | | |
| Table of characteristics: | Present □ | |

| | Not available ⊠ |
|-----------------------------------------------|-----------------|
| Literature: | Present □ |
| (when present, please annex to this document) | Absent ⊠ |