

Lycianthes
Simplified standard protocol: SSP/RTT/1.rev

| | | |
|--|--|-----------------|
| Botanical taxon: | Lycianthes rantonnettii (Carrière) Bitter | |
| Common Name (when known): | Blue Potato Busch | |
| Date of preparation of SSP: | 01-09-2012 | |
| Date of revision of SSP: | 01-01-2020 | |
| SSP revised by: | W.A. Wietsma | |
| Sample to be examined: | VEGETATIVE | |
| Number of growing cycles: | 1 year | |
| Closing date for applications: | 1/12 | |
| Submission date/period: | 1/3 - 31/3 | |
| Seed/Plant Quantity: | 24 young plants | |
| | Able to show all their characteristics in the first year of examination | |
| Special conditions sample: | None | |
| Test station address: | Naktuinbouw, Sotaweg 22, 2371 AA, Roelofarendsveen | |
| Name: | Team Support Variety Testing Department | |
| E-mail: | teamsupport@rasraad.nl | |
| List of grouping characteristics: | NO, (if yes put as annex) | |
| Minimum number of plants in trial: | Vegetative: 20 | Seed: not appl. |
| Minimum number of plants observed by measuring or counting: | Vegetative: 1 | Seed: not appl. |
| Give description of when/where observations on the leaf should take place: | After 6 month's growth | |
| Give description of when/where the other observations should take place: | After 6 month's growth | |
| Test will take place: | GREENHOUSE | |
| Uniformity: | A population standard of 1% with an acceptance probability of at least 95%. Number of Off-types allowed: one off-type allowed in a sample size of 24 | |
| Table of characteristics: | PRESENT (see annex) | |
| | (if present, please annex the table of characteristics and explanations) | |
| Literature: | PRESENT | |
| | (if present, please annex to this document) | |

Table of characteristics Lycianthes

| |
|--|
| Plant: attitude |
| |
| Branches: internodes |
| Branches: shape |
| Branches: juvenile: colour |
| Branches: mature: colour |
| Branches: juvenile: anthocyanin coloration |
| Branches: mature: anthocyanin coloration |
| |
| Leaf: attitude |
| Leaf: petiole: length |
| Leaf: petiole: colour |
| Leaf: petiole: anthocyanin coloration |
| Leaf: blade: shape |
| Leaf: blade: length |
| Leaf: blade: width |
| Leaf: colour of upper side (RHS) |
| Leaf: colour of lower side |
| Leaf: shape top |
| Leaf: shape basal part |
| Leaf: shape in cross-section |
| Leaf: shape margin |
| Young leaves: colour center (RHS) |
| |
| Inflorescence: type |
| |
| Pedice: length |
| Pedice: colour |
| Pedice: anthocyanin coloration at base |
| |
| Calyx: size |
| Calyx: colour |
| Calyx: anthocyanin coloration |
| |
| Corolla: attitude |
| Corolla: shape |
| Corolla: cross-section |
| Corolla: colour (RHS) |
| Corolla: main vein: colour (RHS) |
| Corolla: bullation between veins |
| |
| Stamens: number |
| Stamens: colour |
| |
| Ovary: shape |
| |
| Style |
| |
| Stigma |
| |
| Literature: Dictionary of Gardening: The Royal Horticultural Society Hortica: A.B.Graf The Encyclopedia of Garden Flowers: K&G Bryant |